


Extra questions from Study Unit 7

CHEMIESE EWEWIG. / CHEMICAL EQUILIBRIUM.

Q1


Bepaal K_2 vir die reaksie: / Determine K_2 for the reaction: $\text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \frac{1}{2}\text{O}_2(\text{g})$

as K_1 gelyk is aan $3,4 \times 10^{81}$ by 25°C vir die volgende reaksie: / if K_1 is equal to $3,4 \times 10^{81}$ at 25°C for the following reaction:



Q2

By 900°C is $K_c = 0.0108$ vir die volgende reaksie: / At 900°C , $K_c = 0.0108$ for the following reaction:



Indien 'n mengsel van 15.0 g CaCO_3 , 15.0 g CaO en 4.25 g CO_2 in 'n 10.0 L fles by 900°C geplaas word, sal die hoeveelheid CaCO_3 toeneem, afneem of dieselfde bly wanneer die sisteem ewewig bereik? Omkring die opsie wat jy dink is reg. / If a mixture of 15.0 g CaCO_3 , 15.0 g CaO and 4.25 g CO_2 is placed in a 10.0 L vessel at 900°C , will the amount of CaCO_3 increase, decrease or stay the same when the system reaches equilibrium? Circle the option that you think is correct.









- a) Toeneem. / Increase.
- b) Afneem. / Decrease.
- c) Dieselfde bly. / Stay the same.

Q3

'n Mengsel van CH_4 en H_2O word by 'n 1000 K oor 'n nikkelkatalisator gestuur. Die gasse wat dan vrygestel word, word in 'n 5.00 L fles opgevang. Analise van die gasse by ewewig lewer 0.308 mol CO , 1.287 mol H_2 , 2.681 mol CH_4 en 2.686 mol H_2O . Bereken die ewewigskonstante, K_c , vir die reaksie by 1000 K. / A mixture of CH_4 and H_2O is passed over a nickel catalyst at 1000 K. The emerging gasses is collected in a 5.00 L flask. Analysis of the gasses at equilibrium yielded 0.308 mol CO , 1.287 mol H_2 , 2.681 mol CH_4 and 2.686 mol H_2O . Calculate the equilibrium constant, K_c , for the reaction at 1000 K.

Q4

Die volgende prentjies stel die ewewigstoestand van drie verskillende reaksies voor, van die tipe: / *The following pictures represent the equilibrium state for three different reactions of the type:*




Watter reaksie het die kleinste ewewigskonstante? / *Which reaction has the smallest equilibrium constant?*

Watter reaksie het die grootste ewewigskonstante? / *Which reaction has the largest equilibrium constant?*

Q5

Sal die konsentrasie van NO₂ (g) toeneem, afneem of dieselfde bly as die volgende ewewig versteur word deur:

Will the concentration of NO₂ (g) increase, decrease or remain the same when the following equilibrium is disturbed by:



nog ClNO₂ by te voeg. / *adding more ClNO₂.*

nog ClNO by te voeg. / *adding more ClNO.*

nog NO by te voeg. / *adding more NO.*

NO te verwyder. / *removing NO.*

Q6


Die volgende ewewigskonsentrasies by 127⁰C word waargeneem vir die Haber-proses: / *The following equilibrium concentrations were observed for the Haber process at 127⁰C:*

$$[\text{NH}_3] = 3.1 \times 10^{-2} \text{ mol/L}$$


$$[\text{N}_2] = 8.5 \times 10^{-1} \text{ mol/L}$$

$$[\text{H}_2] = 3.1 \times 10^{-3} \text{ mol/L}$$

- 1 Bereken die waarde van K₁ by 127⁰C vir die reaksie van die Haberproses. / *Calculate the value of K₁ at 127⁰C for the reaction of the Haber process.*
- 2 Bereken die waarde van die ewewigkonstante, K₂, by 127⁰C vir die volgende reaksie.
Calculate the value of the equilibrium constant, K₂, at 127⁰C for the following reaction:



- 3 Bereken die waarde van die ewewigkonstante, K_3 , by 127°C vir die reaksie wat gegee word deur die vergelyking: / Calculate the value of the equilibrium constant, K_3 , at 127°C for the reaction given by the equation:



Q7

Gasagtige NOCl ontbind om die gasse NO en Cl_2 te vorm. By 35°C is die ewewigskonstante 1.6×10^{-5} mol. Wat is die ewewigskonsentrasies wanneer 1.0 mol NOCl in 'n 2.0 liter fles geplaas word?

Gaseous NOCl decomposes to form the gases NO and Cl_2 . At 35°C the equilibrium constant is 1.6×10^{-5} mol/L. When 1.0 mol NOCl is placed in a 2.0 liter flask, what are the equilibrium concentrations?

Q8

Wat is die verskil tussen die ewewigskonstante en die reaksiekwosient? / What is the difference between the equilibrium constant and the reaction quotient?

Q9

Hoekom word gesê dat chemiese ewewigte in 'n dinamiese toestand is? / Why are chemical equilibria said to be in a dynamic state?

- omrede die reagens konsentrasie stadig oor tyd afneem. / because the reactant concentration decreases slowly over time.
- omrede die omgekeerde reaksie enige tyd dominant kan word. / because the reverse reaction could become dominant at any moment.
- omrede die konsentrasies van al die chemiese spesies nie verander nie. / because the concentrations of all chemical species do not change.
- omrede die voorwaartse en terugwaartse reaksies teen dieselfde tempo aanhou. / because the forward and reverse reactions continue to occur at equal rates.

Q10

0.55 M COI_2 word aanvanklik in 'n 2.00 L fles verhit en die reaksie word toegelaat om ewewig te bereik. By ewewig is die konsentrasie I_2 gelyk aan 0.022 M. Bereken die ewewigskonstante, K_c , vir die reaksie.

0.55 M of COI_2 is heated initially in a 2.00 L flask and the reaction is left to reach equilibrium. At equilibrium the concentration of I_2 is equal to 0.022 M. Calculate the equilibrium constant, K_c , for the reaction.

